## Problem assignment 1.

## Meromorphic Continuation of Eisenstein Series.

Joseph Bernstein

July 8, 2007.

I would like to discuss the notion of **Frechet** representation, as discussed in lectures (Casselman uses the term "Frechet representations of moderate growth").

Bellow I will consider the group  $G = SL(2, \mathbf{R})$ . For  $g \in G$  I define  $||g|| = max(||g||_M, ||g^{-1}||_M)$ , where  $||g||_M$  is just the usual norm on the space of matrices.

**1.** Let C(G) be the space of continuous functions on G. Show that this is a Frechet space in a natural topology, that the natural representation  $\Pi, G, C(G)$  is continuous, but it is not a Frechet representation.

**2.** Prove the following

**Lemma.** Let A be a \*-algebra and  $\pi : A \to Op(H)$  its \*-representation in a Hilbert space H. Suppose we know that the subalgebra  $B \subset Op(H)$  contains many compact operators (which means that they do not have common kernel in H).

Show that the representation  $\Pi$  is isomorphic to a sum of irreducible representations  $\Pi = \oplus \pi_{\kappa}$ , and each irreducible representation appears in this decomposition with finite multiplicity.

**3.** Let *M* be a  $(\mathfrak{g}, K)$ -module. Suppose we know that for any *K*-type  $\sigma$  the space  $M^{\sigma}$  is finite dimensional.

Show that the following conditions are equivalent

(i) M is finitely generated

(ii) M is  $\mathcal{Z}(G)$  finite

(iii) M has finite length.

(here  $\mathcal{Z}(G)$  is the center of the Universal enveloping algebra  $U(\mathfrak{g})$ ) Such modules are called **Harish Chandra modules**.

4. Using Casselman-Wallach theorem prove the following

**Statemant.** Let  $(\pi, G, V)$  be an asf representation (i.e. an admissible smooth Frechet representation). Then for any Frechet representation ((R, G, E) we have

$$Hom_G(V, E) = Hom_G(V, E^{\infty}) = Hom_{\mathcal{H}^f}(V^f, E^{\infty f}) = Hom_{(\mathfrak{g}, K)}(V^f, E^{\infty f})$$

c c

c

.

5. Prove Frobenius reciprocity.

**Statement.** Let  $(\pi, G, V)$  be an asf representation,  $X = \Gamma \backslash G$  an automorphic space. Show that

$$Hom_G(V, F(X)) = Hom_G(V, C^{\infty}(X)) = Hom_{\Gamma}(V, \mathbf{C})$$

6. Let  $(\pi, G, V)$  be an asf representation. Define the notion of a contra-gradient asf representation  $\tilde{V} \subset V^*$  and of hermitian dual asf representation  $V^+ = \bar{\tilde{V}}.$ 

Show that for any Banach representation (R,G,E) and a  $G\operatorname{-moprphism}\nu$  :  $V \to E$  the adjoint morphism  $\nu^*$  maps  $(E^*)^{\infty}$  into  $\tilde{V}$ . Show that morphisms  $\alpha : V \to C^{\infty}(X)$  correspond to morphisms  $\beta$ :

 $C_c^{\infty}(X) \to V^+.$